

# Calculus I - Lecture 6

## Limits D & Intermediate Value Theorem

Lecture Notes:

<http://www.math.ksu.edu/~gerald/math220d/>

Course Syllabus:


<http://www.math.ksu.edu/math220/spring-2014/indexs14.html>

Gerald Hoehn (based on notes by T. Cochran)


February 5, 2014

## Section 2.7 – Limits at Infinity

**Recall:** 1) A **vertical asymptote** is a guideline that the graph of  $f(x)$  approaches at points where  $\lim_{x \rightarrow a^+} f(x) = \pm\infty$  or  $\lim_{x \rightarrow a^-} f(x) = \pm\infty$ .



2) A **horizontal asymptote** is a guideline that the graph of  $f(x)$  approaches at points where  $x \rightarrow \pm\infty$ .



# Limits at Infinity

## Definition

The limits of a function  $f(x)$  **at infinity** are the values  $f(x)$  approaches as  $x \rightarrow \infty$ , written  $\lim_{x \rightarrow \infty} f(x)$ , or  $x \rightarrow -\infty$ , written  $\lim_{x \rightarrow -\infty} f(x)$ . They may or may not exist.

**Example:** In the previous example we find:

$$\lim_{x \rightarrow \infty} f(x) = -3 \quad (\text{horizontal asymptote to the right})$$

$$\lim_{x \rightarrow -\infty} f(x) = 2 \quad (\text{horizontal asymptote to the left})$$

**Note:**

$$\lim_{x \rightarrow \infty} f(x) = L \Leftrightarrow y = L \text{ horizontal asymptote for } x \rightarrow \infty$$

$$\lim_{x \rightarrow -\infty} f(x) = L \Leftrightarrow y = L \text{ horizontal asymptote for } x \rightarrow -\infty$$

$\frac{\infty}{\infty}$ -type limits (Example:  $\lim_{x \rightarrow \infty} \frac{x+2}{x-3}$ )

**Basic Trick** for evaluating  $\frac{\infty}{\infty}$ -type limits (without doing any graphing!):

Divide top and bottom by the largest power of  $x$  occurring in the denominator.

**Example:** a) Evaluate  $\lim_{x \rightarrow \infty} \frac{2x + 3}{2 - x}$ .

**Solution:**  $x$  appears with first degree in denominator.

$$\begin{aligned}\lim_{x \rightarrow \infty} \frac{2x + 3}{2 - x} &= \lim_{x \rightarrow \infty} \frac{(2x + 3) \cdot \frac{1}{x}}{(2 - x) \cdot \frac{1}{x}} \\ &= \lim_{x \rightarrow \infty} \frac{2x \cdot \frac{1}{x} + 3 \cdot \frac{1}{x}}{2 \cdot \frac{1}{x} - x \cdot \frac{1}{x}} \\ &= \lim_{x \rightarrow \infty} \frac{2 + \frac{3}{x}}{\frac{2}{x} - 1} \\ &= \frac{\lim_{x \rightarrow \infty} (2 + \frac{3}{x})}{\lim_{x \rightarrow \infty} (\frac{2}{x} - 1)} = \frac{2 + 0}{0 - 1} = -2\end{aligned}$$

b) What information does the limit in part a) provide about the graph of  $f(x) = \frac{2x+3}{2-x}$ ?

**Solution:**

Horizontal asymptote:  $y = -2$  (for  $x \rightarrow \infty$ )

$$\lim_{x \rightarrow \infty} \frac{2x+3}{2-x} = -2 \quad (\text{by same method as in part a)})$$

Horizontal asymptote:  $y = -2$  (for  $x \rightarrow -\infty$ )

## Rational Functions

**Example:** Evaluate  $\lim_{x \rightarrow \infty} \frac{2x - 5x^3}{x^3 - x + 1}$ .

**Solution:**  $x$  appears with 3<sup>rd</sup> degree in denominator.

$$\begin{aligned}\lim_{x \rightarrow \infty} \frac{2x - 5x^3}{x^3 - x + 1} &= \lim_{x \rightarrow \infty} \frac{(2x - 5x^3) \frac{1}{x^3}}{(x^3 - x + 1) \frac{1}{x^3}} \\ &= \lim_{x \rightarrow \infty} \frac{\frac{2x}{x^3} - \frac{5x^3}{x^3}}{\frac{x^3}{x^3} - \frac{x}{x^3} + \frac{1}{x^3}} \\ &= \lim_{x \rightarrow \infty} \frac{\frac{2}{x^2} - 5}{1 - \frac{1}{x^2} + \frac{1}{x^3}} = \frac{0 - 5}{1 - 0 + 0} = -5\end{aligned}$$

## Alternate way:

This works for any rational function (quotient of polynomials)

$$\lim_{x \rightarrow \infty} \frac{a x^n + \text{lower degree terms}}{b x^m + \text{lower degree terms}} = \lim_{x \rightarrow \infty} \frac{a x^n}{b x^m}.$$

Drop all lower degree terms.

**Example:** Redo the limit  $\lim_{x \rightarrow \infty} \frac{2x - 5x^3}{x^3 - x + 1}$  using the alternate way.

## Solution:

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{2x - 5x^3}{x^3 - x + 1} &= \lim_{x \rightarrow \infty} \frac{-5x^3 \overset{\text{lower}}{\overbrace{+ 2x}}}{x^3 \underbrace{-x + 1}_{\text{lower}}} \\ &= \lim_{x \rightarrow \infty} \frac{-5x^3}{x^3} \\ &= \lim_{x \rightarrow \infty} \frac{-5}{1} = -5 \quad (\text{Books method}) \end{aligned}$$

**Example:** Evaluate  $\lim_{x \rightarrow \infty} \frac{5x^4 - 2x}{6x^3 + 7x^5 - 3}$ .

**Solution:**

**First method:**  $x$  appears with 5<sup>th</sup> degree in denominator.

$$= \lim_{x \rightarrow \infty} \frac{(5x^4 - 2x) \frac{1}{x^5}}{(6x^3 + 7x^5 - 3) \frac{1}{x^5}}$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{5}{x} - \frac{2}{x^4}}{\frac{6}{x^2} + 7 - \frac{3}{x^5}}$$

$$= \frac{0 - 0}{0 + 7 - 0} = \frac{0}{7} = 0$$

**Second method:**

$$= \lim_{x \rightarrow \infty} \frac{5x^4}{7x^5}$$

$$= \lim_{x \rightarrow \infty} \frac{5}{7x} = 0$$

## $\frac{\infty}{\infty}$ type limits with radicals

**Example:** Compute  $\lim_{x \rightarrow -\infty} \frac{\sqrt{4x^2 - 2}}{x + 3}$ .

**Solution:**

$$\begin{aligned}\lim_{x \rightarrow -\infty} \frac{\sqrt{4x^2 - 2}}{x + 3} &= \lim_{x \rightarrow -\infty} \frac{\sqrt{4x^2 - 2} \cdot \frac{1}{x}}{(x + 3) \cdot \frac{1}{x}} \\ &= \lim_{x \rightarrow -\infty} \frac{\sqrt{x^2(4 - \frac{2}{x^2})} \cdot \frac{1}{x}}{1 + \frac{3}{x}}\end{aligned}$$


$\sqrt{x^2} = x$ ?? One has:  $\sqrt{x^2} = x$  if  $x > 0$  and  $\sqrt{x^2} = -x$  if  $x < 0$ .  
because  $x < 0$

$$\begin{aligned}&= \lim_{x \rightarrow -\infty} \frac{\overbrace{-x} \sqrt{4 - \frac{2}{x^2}} \cdot \frac{1}{x}}{1 + \frac{3}{x}} \\ &= \lim_{x \rightarrow -\infty} \frac{-\sqrt{4 - \frac{2}{x^2}}}{1 + \frac{3}{x}} \\ &= \frac{-\sqrt{4 - 0}}{1 + 0} = \frac{-2}{1} = -2\end{aligned}$$

## Section 2.8 – Intermediate Value Theorem

### Theorem (Intermediate Value Theorem (IVT))

Let  $f(x)$  be **continuous** on the interval  $[a, b]$  with  $f(a) = A$  and  $f(b) = B$ .




*Given any value  $C$  between  $A$  and  $B$ , there is at least one point  $c \in [a, b]$  with  $f(c) = C$ .*

**Example:** Show that  $f(x) = x^2$  takes on the value 8 for some  $x$  between 2 and 3.

**Solution:** One has  $f(2) = 4$  and  $f(3) = 9$ . Also  $4 < 8 < 9$ . Since  $f(x)$  is continuous, by the IVT there is a point  $c$ , with  $2 < c < 3$  with  $f(c) = 8$ .

**Note:** The IVT fails if  $f(x)$  is not continuous on  $[a, b]$ .


**Example:**



There is no  $c \in [a, b]$  with  $f(c) = C$ .

### Important special case of the IVT:

Suppose that  $f(x)$  is **continuous** on the interval  $[a, b]$  with  $f(a) < 0$  and  $f(b) > 0$ .



Then there is a point  $c \in [a, b]$  where  $f(c) = 0$ .

**Example:** Show that the equation

$$x^3 - 3x^2 + 1 = 0$$

has a solution on the interval  $(0, 1)$ .

**Solution:**

- 1)  $f(x) = x^3 - 3x^2 + 1$  is continuous on  $[0, 1]$  (polynomial).
- 2)  $f(0) = 1 > 0$ .
- 3)  $f(1) = 1 - 3 + 1 = -1 < 0$ .

Thus by the IVT, there is a  $c \in (0, 1)$  with  $f(c) = 0$ .